

ROSAT X-RAY OBSERVATIONS OF CENTRAL GALACTIC STARBURSTS

D. Tschöke and G. Hensler

Institute of Theoretical Physics and Astrophysics, University of Kiel, Germany

RESUMEN

El resumen será traducido al español por los editores.

We present spatial and spectral results of ROSAT observations of two galaxies: NGC 4569 and NGC 4303. Both objects are members of the Virgo Cluster and have nearby smaller companions with which they have probably gone through an encounter.

Both soft X-ray spectra (0.1-2.4 keV) can be fitted with a two-component model consisting of a power-law component from a non-thermal emission of a compact active nucleus and a Raymond-Smith model for a hot thin thermal plasma originating from starbursts. The X-ray luminosities are in the range of 10^{40} erg s⁻¹. The central source of NGC 4569 is very compact and can not be resolved with the ROSAT HRI detector, while there is extended soft X-ray emission in the galactic disk as well as above it. NGC 4303 is dominated by the active nucleus in the X-ray. But several additional disk sources associated with HII regions can be observed.

ABSTRACT

We present spatial and spectral results of ROSAT observations of two galaxies: NGC 4569 and NGC 4303. Both objects are members of the Virgo Cluster and have nearby smaller companions with which they have probably gone through an encounter.

Both soft X-ray spectra (0.1-2.4 keV) can be fitted with a two-component model consisting of a power-law component from a non-thermal emission of a compact active nucleus and a Raymond-Smith model for a hot thin thermal plasma originating from starbursts. The X-ray luminosities are in the range of 10^{40} erg s⁻¹. The central source of NGC 4569 is very compact and can not be resolved with the ROSAT HRI detector, while there is extended soft X-ray emission in the galactic disk as well as above it. NGC 4303 is dominated by the active nucleus in the X-ray. But several additional disk sources associated with HII regions can be observed.

Key Words: X-RAYS: GALAXIES — GALAXIES: ACTIVE — GALAXIES: STARBURSTS — GALAXIES: INDIVIDUELL (NGC 4303, NGC 4569)

1. INTRODUCTION

The phenomenon of galactic nuclear activity is strongly connected with starbursts, especially in galaxies with low luminous active nuclei, like LINERs and Seyfert galaxies. Beside UV and optical observations the analysis of X-ray data reveals composite spectra, indicating non-thermal nuclear activity and soft thermal emission from the hot gas of a central starburst. One main aspect of interest is the triggering mechanism of galactic activity in different morphologies, like interacting systems (Tschöke et al. 1999) and bars.

Here we investigate luminosity and spectral distribution of the soft X-ray emission from two nearby barred galaxies with nuclear starburst activity. In both nearby galaxies the X-ray emission can be clearly distinguished between different galactic components: the center, the disk, and, in the case of NGC 4569, the halo. Both galaxies have been classified as LINERS from optical spectroscopy (, NGC 4569: Willner et al. 1985; NGC 4303: Huchra et al. 1982).

NGC 4569 is a bright early-type spiral in the Virgo Cluster, one of the few blue-shifted galaxies outside the local group. It is gas-deficient in the outer spiral arms, the neutral hydrogen content is strongly concentrated in the inner region (Cayatte et al. 1990). The bright nucleus, embedded in a normal stellar bulge, is probably the result of a recent star formation episode (Stauffer et al. 1986).

NGC 4303 is a barred late-type spiral galaxy and also member of the Virgo Cluster. Indications for a high star formation rate in NGC 4303 are the high density of HII regions (Hodge & Kennicutt 1983; Martin & Roy 1992) and three observed supernovae (van Dyk 1992). It shows also strong radio emission distributed over the entire disk (Condon 1983). Two nearby galaxies, NGC 4303 A (Condon 1983) and NGC 4292 (Cayatte et al. 1990), are possible candidates for a past interaction with NGC 4303.

2. SPATIAL DISTRIBUTION, X-RAY SPECTRA AND LUMINOSITIES

2.1. NGC 4569

The global X-ray spectrum of NGC 4569 can be described by a two-component fit consisting of a power-law component (predominantly from the nucleus) and a thermal plasma component (hot gas from SNRs and superbubbles). The resulting value of hydrogen absorption is in agreement with the Galactic foreground, supporting the observed HI deficiency (Cayatte et al. 1990). The total luminosity in the ROSAT band for NGC 4569 amounts to $\sim 2 \times 10^{40}$ erg s $^{-1}$.

Two interpretations for the emission of the compact nucleus in the HRI image seem possible: either X-ray emission from an unresolved compact nuclear source, to be associated with the LINER-type nucleus of NGC 4569 in the optical, or a slightly extended nuclear starburst region, similar to the one found in NGC 1808 (Junkes et al. 1995) at a scale of ~ 1 kpc. The spectrum of the nuclear source can well be fitted with a power law spectrum - with the spectral resolution of the ROSAT PSPC it is not possible to clearly determine the nature of the central source - either the contribution from high-mass binaries in the central starburst, or an AGN component with a spectral distribution in the ROSAT band which could be described by a power-law fit.

X-ray emission from the disk of NGC 4569 is clearly visible in the PSPC image (Fig. 1). The total disk emission in X-rays from either side (NE vs. SW) is comparable, Their spectral energy distribution however differ significantly. The contribution from the northern part is significantly harder. This may indicate either higher absorption in the north, or an intrinsically harder X-ray spectrum caused by the cumulative impact of several SNRs or high-mass X-ray binaries. Higher internal absorption towards the northern part of NGC 4569 seems difficult to explain since the HI content of that galaxy is strongly confined to the central part (Cayatte et al. 1990).

NGC 4569 has an inclination of $\sim 65^\circ$. The galaxy appears extended in soft X-rays, the X-ray emission in the ROSAT band shows contributions from areas outside the nucleus and outside the disk. In contrast to edge-on galaxies, where emission components from disk and halo can be quite easily distinguished, it is more difficult to disentangle these components in the case of NGC 4569.

The overall distribution of the soft X-ray emission appears to be very asymmetric with a spur extending from the galaxy center and some additional diffuse patches west of the galaxy. The source northwest of the galactic center is probably not connected to NGC 4569. A further source southwest of the nucleus of NGC 4569 can be identified with a background galaxy visible in the optical. We derived a hardness ratio (0.5-2.0 keV vs. 0.1-0.4 keV) of -0.02 ± 0.17 for the area west of NGC 4569, excluding both these independent X-ray sources. The total number of counts after background subtraction for this area is only 68, hence no meaningful spectrum could be derived. The hardness ratio is compared with the results from model spectra. A value of $HR = 0.0$ results for a thermal plasma spectrum (e.g. Raymond & Smith 1977) with a plasma temperature of 0.2 keV. Hence, the X-ray emission from the area outside the disk can be explained with hot gas of a temperature of approximately 2×10^6 K.

An additional argument in favour of diffuse X-ray emission in the halo of NGC 4569 comes from the very

good spatial correlation of the soft X-ray (0.1-0.4 keV) emission with a large diffuse H α spur (Fig. 1). Similar correlations were observed in a number of nearby starburst galaxies (e.g. M82: Shopbell & Bland-Hawthorn 1998, NGC 3628: Dahlem et al. 1996, NGC 4666: Dahlem et al. 1997) and are interpreted as a signature for an outflow of hot gas from the starburst center. Our X-ray data are fully consistent with such an interpretation in the case of NGC 4569.

2.2. NGC 4303

Obvious differences between NGC 4303 and NGC 4569 are that (1) NGC 4303 has a much lower inclination (27° ; Guhathakurta et al. 1988) and (2) the amount of HI is more distributed over the total galactic disk and much higher in the outer parts than in NGC 4569, which is closer to the Virgo Cluster center. Consequently, the first point allows better observation possibilities to distinguish between disk and central source. Vice versa, it is not possible to observe a soft halo component above the galactic disk, as can be seen in NGC 4569 or in more details in edge-on starburst galaxies (e.g. Vogler et al. 1995).

Another difficulty in the X-ray observations of NGC 4303 is that no spatial distribution can be obtained from the PSPC data due to the fact that the galaxy is $17'$ off-axis in the field of view. This degenerates the spatial resolution of the observation to $\sim 67''$.

The global X-ray spectrum of NGC 4303 can be described by a similar fit to that of NGC 4569 with a non-thermal power-law component and a Raymond-Smith component. The resulting value of hydrogen absorption is with $3.4 \times 10^{20} \text{ cm}^{-2}$ by a factor of 2 higher than the Galactic foreground. This is in agreement with the distribution of HI gas over the whole galactic disk of NGC 4303. The total luminosity of NGC 4303 in the ROSAT band is $\sim 5 \times 10^{40} \text{ erg s}^{-1}$.

The HRI image reveals a dominant central source and several additional sources distributed along the spiral arms (Fig. 2). UV observations (Colina & Arribas 1999) result in an unresolved ($\leq 8 \text{ pc}$) active nucleus (compact stellar cluster or AGN) and a surrounding spiral-shaped region of very young massive star forming clusters (2-3 Myr). This picture leads to attribute a power-law component to the nuclear source, contributing more than 80% to the total spectrum. This is in agreement with the results from the HRI observation, where the main part originates from the center of the galaxy. The coincidence of the disk sources with HII regions in the spiral arms is in favour of several star forming regions with SNRs to be the origin of the observed X-ray emission. The sources at the ends of the bar are in agreement with expected gas flows and accumulation within the bar. The eastern spiral feature in the optical and H α with the X-ray source at its bend suggest a past tidal disturbance in the gas dynamics leading to enhanced star formation.

Further contribution to the X-ray flux from HMXBs in the disk sources and from a circumnuclear starburst at a distance up to $\sim 1 \text{ kpc}$ from the center can not be excluded from the HRI observation.

3. CONCLUSIONS

A comparison of the total luminosities of NGC 4569 and NGC 4303 in the soft X-ray with a larger sample of normal and active galaxies (Green et al. 1992) shows good agreement with the galaxies (including LINERs and starburst galaxies) in that sample. Most of the soft X-ray emission in both galaxies comes from their nuclear regions, but additional components from the disk and, in the case of NGC 4569, possibly from the halo, can be distinguished. The existence of a central superwind in NGC 4569 is supported by two facts: First, the extended emission component is predominantly at one side coincident with an H α feature at high z . This one-sided geometry would be expected in an ambient medium with a density gradient along the z -direction and a starburst region slightly above the galactic plane. Second, its spectral distribution (hardness ratio) can be explained with a hot plasma of a temperature of a few 10^6 K , which would be expected in the halo of starburst galaxies.

The nature of the nuclei in NGC 4569 and NGC 4303 is still unknown. X-ray images and spectra indicate a coexistence of a dominating AGN and a starburst in both galaxies. Both harbour a compact central source, emitting most of the soft X-ray. Nevertheless, massive nuclear stellar clusters with enhanced star formation can not be excluded to account for the sources.




Figure 1. NGC 4569: PSPC X-ray contours over $H\alpha$ image.




Figure 2. NGC 4303: HRI X-ray contours over R band image (Frei et al. 1996).

REFERENCES

- Cayatte, V., van Gorkom, J. H., Balkowski, C. & Kotanyi, C. 1990, AJ 100, 604
Colina, L. & Arribas, S. 1999, ApJ 514, 637
Condon, J. J. 1983, ApJS 53, 459
Dahlem, M., Heckman, T. M., Fabbiano, G., Lehnert, M. D. & Gilmore, D. 1996, ApJ 461, 724
Dahlem, M., Petr, M. G., Lehnert, M. D., Heckman, T. M. & Ehle, M. 1997, A&A 320, 731
Frei, Z., Guharthakurta, P., Gunn, J. E. & Tyson J. A. 1996, AJ 111, 174
Green, P. J., Anderson, S. F. & Ward, M. J. 1992, MNRAS 254, 30
Guhathakurta, P., van Gorkom, J. H., Kotanyi, C. G. & Balkowski, C. 1988, AJ 96, 851
Hodge, P. W. & Kennicutt, R. C. 1983, AJ 88, 296
Huchra, J. P., Wyatt, W. F. & Davies, M. 1982, AJ 87, 1628
Junkes, N., Zinnecker, H., Hensler, G., Dahlem, M. & Pietsch, W. 1995, A&A 294, 8
Martin, P. & Roy, J.-R. 1992, ApJ 397, 463
Raymond, J. C. & Smith, B. W. 1977, ApJS 35, 419
Shopbell, P. L. & Bland-Hawthorn, J. 1998, ApJ 493, 129
Stauffer, J. R., Kenney, J. D. & Young, J. S. 1986, AJ 91, 1286
Tschöke, D., Hensler, G. & Junkes, N. 1999, A&A 343, 373
van Dyk, S. D. 1992, AJ 103, 1788
Vogler, A., Pietsch, W. & Kahabka, P. 1995, Advances in Space Research, vol. 16, no. 3, p. 139
Willner, S. P., Elvis, M. & Fabbiano, G. 1985, ApJ 299, 443

D. Tschöke and G. Hensler: Institute of Theoretical Physics and Astrophysics, University of Kiel, Leibnizstr. 15, D-24098 Kiel, Germany (tschoeke,hensler@astrophysik.uni-kiel.de)